


Seasonal weather forecast for the months of June, July and August 2017

General over view of the weather expected to prevail for June, July and August.

The weather for the following three months, namely **June, July and August 2017**, will be presented, with respect to the anomalies of temperature, and accumulated precipitation.

Normal temperatures (both maximum and minimum), are on the rise suggesting the second half of the period as the hottest period of the year. The accumulated precipitation is at its lowest values. The period of the following three months is likely to be characterized by outbreaks of thunderstorms, initiated from thermal causes over the main plain area or over the mountainous range from orographic uplift.

Specifically, regarding the seasonal forecast for the period of **June, July and August 2017**, the temperature will be generally normal. Normal temperature characterizes also the greater part of Asia Minor, while the Balkans will be warmer than normal and near East will be lower than normal. The amount of the accumulated precipitation over Cyprus is expected to range generally below 50% of normal as the Balkans and part of Asia Minor, except the northern part where the accumulated precipitation will range between 50% and 75% of normal.




Divergence of temperature from normal from the mean seasonal temperature (°C) for June, July and August

Percentage (%) of the mean seasonal accumulated precipitation compared to the mean normal for June, July and August

The seasonal monthly forecast in detail

June

The seasonal forecast suggests that **June's** temperature will be generally normal while the Balkans and great parts of Asia Minor will be warmer than normal. Regarding the accumulated precipitation Cyprus is expected to be dry with the accumulated precipitation ranging below 50% for the entire area of Cyprus.




Divergence of the mean monthly temperature (°C)
from normal during June


Percentage of the mean monthly precipitation (%)
compared with the normal of June

July

The seasonal forecast suggests that **July's** temperature will be over the greater part of the island normal but cooler by 1 to 2°C, over the extreme east part. Regarding the accumulated precipitation Cyprus west part is expected to be again with the accumulated precipitation ranging below 50% but over the east part accumulated precipitation of up to 200% is probable, bearing always in mind that the normal accumulation for July is very low.




Divergence of the mean monthly temperature (°C) from normal during July




Percentage of the mean monthly precipitation (%) compared with the normal of July

August

The seasonal forecast suggests that **August's** temperature will be generally below normal, by 1 to 2°C. Regarding the accumulated precipitation Cyprus will be again with the accumulated precipitation ranging between 50% to 75% of normal, except southeastern area where the accumulated precipitation will be below 50% of normal.



Divergence of the mean monthly temperature (°C) from normal during August



Percentage of the mean monthly precipitation (%) compared with the normal of August

Normal values of temperature (mean maximum and mean minimum) and accumulated precipitation for June, July and August

The normal values of mean maximum, mean minimum temperature and accumulated precipitation are presented below concerning the three months the period of forecast is covering, in order to gain a better view of the normal seasonal climate. The period of forecast is the transitional summer. The temperature, both maximum and minimum, is on the rise with **July's** temperature significantly higher. Climatologically extreme high temperatures are recorded during **July** and over some areas during **August**. On the other hand the accumulated precipitation is declining, as presented by the climatological data shown on the table below. Thundery activity is likely during afternoon hours mainly inland and over the mountains, which is noted also from the table below. Note the cause of the mountainous area of Troodos range over the accumulated precipitation, mainly as a result of thundery activity, which is affecting primarily the mountainous ranges of the island. The accumulated precipitation over Prodromos station although declining towards **August** is always significantly higher, if compared to the accumulated precipitation of the other selected stations.


TEMPERATURE AND PRECIPITATION NORMAL VALUES FOR THE PERIOD 1981-2010									
Area Name	MEAN DAILY MAXIMUM TEMPERATURE (°C)			MEAN DAILY MINIMUM TEMPERATURE (°C)			MEAN MONTHLY TOTAL PRECIPITATION (mm)		
	June	July	August	June	July	August	June	July	August
NORTH COAST	30.3	33.4	33.3	18.5	21.1	21.5	1.8	0.1	0.0
WEST COAST*	27.6	29.9	30.5	18.0	20.6	21.1	1.3	0.2	0.0
MOUNTAINOUS AREAS	29.0	27.9	28.0	15.0	18.2	18.1	27.2	16.4	12.0
INLAND*	34.0	37.1	36.9	19.1	22.1	21.9	11.6	4.2	1.8
SOUTH COAST	30.2	32.5	32.9	19.4	22.0	22.3	2.0	0.5	0.3
EAST COAST**	31.0	33.2	33.5	18.9	21.8	22.0	1.8	0.1	0.1

* West Coast and Inland Values cover the period 1983-2010

** East Coast Temperature Values cover the period 1981-2007

Evaluation of Mays seasonal forecast for the area of Cyprus

The seasonal forecast suggested that **May's** temperature would be generally normal while normal to slightly above normal was suggested to be over most parts of the surrounding area. Regarding the accumulated precipitation was expected to range below 50% of normal over south and east areas and below 75% over the rest of the island.



Divergence of the mean monthly temperature (°C)
from normal during May

Percentage of the mean monthly precipitation (%)
compared with the normal of May

Concerning the models evaluation for **May**, a general judgment is that the model performed relatively well concerning only the temperature but it did not perform so well in forecasting the seasonal accumulated precipitation.

In a more detailed evaluation of the recorded temperatures both maximum and minimum (table below) were around normal, except Prodromos station where both mean daily maximum temperature and mean daily minimum temperature were above normals. Great departures were also recorded in several cases except Prodromos station, such as the maximum of Polis Chrysochous station that was 33.7°C (normal 25.8°C), and also the maximum of Larnaca airport station that was 34.2°C (normal 26.4°C). A daily maximum temperature of 24.2°C (with a normal of 29.6°C) that was well below normal was recorded at Athalassa station.

Daily minimum temperatures that were above normal (deviating by 4°C or more from normal) were also recorded in many stations. Except the highest daily minimum temperature of Prodromos that was 21.6°C (with a normal of 10.8°C), note the highest daily minimum of Athalassa that was 20.5°C (with a normal of 14.8°C).

From the distribution (provisional accumulated precipitation chart) of the accumulated precipitation of the month the mean surface distribution was near normal (reaching 15.6mm or 98% of normal), and not as low as the model suggested. It is obvious, that the model did not perform so well concerning the accumulated precipitation.


It is worth mentioning that from the provisional data on the 2nd, 3rd, 5th and 23rd of the month hail was reported.

TEMPERATURE AND PRECIPITATION PROVISIONAL DATA FOR MAY 2017														
St. No.	Station Name	Mean Daily Maximum Temperature (°C)	Normal Value (1981-2010)	Difference from Normal Value	Highest Daily Maximum Temperature (°C)	Lowest Daily Maximum Temperature (°C)	Mean Daily Minimum Temperature (°C)	Normal Value (1981-2010)	Difference from Normal Value	Lowest Daily Minimum Temperature (°C)	Highest Daily Minimum Temperature (°C)	Monthly Total Precipitation (mm)	Normal Value (1981-2010)	Difference from Normal Value
41	POLIS CHRYSOCHOUS	26.0	25.8	0.2	33.7	22.5	14.5	14.3	0.2	12.0	18.7	9.1	7.6	1.5
82*	PAFOS (AIRPORT)	25.3	24.5	0.8	29.7	22.4	15.0	14.7	0.3	12.0	19.2	13.0	6.1	6.9
225	PRODROMOS (C.F.C.)	23.6	20.4	3.2	33.4	11.0	12.5	10.8	1.7	7.9	21.6	109.1	32.0	77.1
666*	ATHALASSA (RADIOSONDE)	29.3	29.6	-0.3	36.6	24.2	15.6	14.8	0.8	12.7	20.5	5.6	24.6	-19.0
731	LARNAKA (AIRPORT)	26.9	26.4	0.5	34.2	23.8	16.2	15.7	0.5	13.6	20.2	27.6	9.8	17.8
800**	ACHNA (DASAKI)	27.2	27.4	-0.2	33.9	23.5	14.8	14.9	-0.1	12.2	19.7	24.2	10.7	13.5

* Pafos' and Athalassa's Station Normal Values cover the period 1983-2010
** Achna's Temperature Normal Values cover the period 1981-2007

= VALUES FROM AUTOMATIC WEATHER STATION

For the purpose of better visualization of **May** accumulated precipitation, a chart of Cyprus with the total preliminary accumulated precipitation is presented.



A Cyprus chart with the normal (period 1981 to 2010) accumulated precipitation for the month of **May** is also presented.

