Seasonal weather forecast for the months of


December 2020 and January and February 2021

General over view of the weather expected to prevail during December 2020 and January and February 2021.

The following period consists of the season of the traditional winter, where most of the annual accumulated precipitation is climatologically expected while in the period in focus normal temperatures (both maximum and minimum) are significantly lower than the other months of the year. During the period, the frequency of baroclinic depressions affecting the area is at its most resulting in higher accumulated precipitation amounts.

Seasonal forecast for the next three months

Specifically, regarding the seasonal forecast for the period of **December 2020** and **January** and **February 2021**, the temperature is expected to be generally slightly above normal. Normal to above normal temperature characterizes also Asia Minor and the Balkans. The amount of the accumulated precipitation inland and over the southern and eastern part of Cyprus is expected to range generally from 50% to 75% of normal while over the southeast part will range below 50% of normal (*).




Divergence of temperature from normal from the mean seasonal temperature (°C) for December, January and February Percentage (%) of the mean seasonal accumulated precipitation compared to the mean normal for December, January and February


Seasonal forecast in detail

The seasonal forecast for **December 2020** suggests that temperature will be above normal (by 1°C to 2°C) mainly over coastal areas. The accumulated precipitation of **December** is forecasted to be disappointing since it suggests a mainly dry month with accumulation ranging only from 50% to 75% of

normal, except the south eastern part where the accumulation will range below 50% of normal. Low accumulations characterize all the surrounding area (*).




The seasonal forecast for **January 2021** suggests that temperature will be above normal (by 1°C to 2°C). The forecast of **January** for the accumulated precipitation suggests that the accumulation will range between from 50% to 75% of normal all over the island, except the south eastern coastal parts where the accumulation will range below 50% of normal. Low accumulations characterize almost all the surrounding area (*).



The seasonal forecast for **February 2021** is almost similar to the one of January and it will be warmer than normal by 1°C to 2°C. The accumulated precipitation of **February** forecast suggests that the

accumulation will range between from 50% to 75% of normal all over the island, except the south eastern parts where the accumulation will range below 50% of normal. Low accumulations are characterizing great parts of the surrounding area (*).



Divergence of the mean monthly temperature (°C) from normal during February

Percentage of the mean monthly precipitation (% compared with the normal of February

Normal values of temperature (mean maximum and mean minimum) and accumulated precipitation for December 2020 and January and February 2021

The normal values of mean maximum, mean minimum temperature and accumulated precipitation are presented below concerning the three months the period of forecast is covering, in order to gain a better view of the normal seasonal climate. The temperatures, both maximum and minimum, are the lowest climatological temperatures of the year while the accumulated precipitation has the greater amount of the year, because of the increased frequency of reoccurrence/redeveloping of depressions over the area of the east Mediterranean. During the period of forecast, snow is likely to occur over the Troodos range while, in the event of the likely synoptic situation, snow may occur above 300m a phenomenon not that frequent.


| | TEM | PERATURE | AND PREC | I NOITATION | NORMAL V | ALUES FOR | THE PERIO | D 1981-20 | 10 | |
|-------------------|----------|------------------------|----------|--|----------|-----------|---------------------------------------|-----------|----------|--|
| | | DAILY MAX 1PERATURE | | MEAN DAILY MINIMUM TEMPERATURE (°C) | | | MEAN MONTHLY TOTAL PRECIPITATION (mm) | | | |
| Area Name | December | January | February | December | January | February | December | January | February | |
| NORTH COAST | 18.0 | 16.3 | 16.3 | 9.3 | 7.6 | 7.3 | 93.6 | 85.3 | 68.5 | |
| WEST COAST* | 18.9 | 17.1 | 17.1 | 10.0 | 8.3 | 8.1 | 90.1 | 78.8 | 59.8 | |
| MOUNTAINOUS AREAS | 8.3 | 6.3 | 6.7 | 2.6 | 0.7 | 0.5 | 157.3 | 150.0 | 128.7 | |
| INLAND* | 17.3 | 15.5 | 16.0 | 7.0 | 5.4 | 5.3 | 57.2 | 48.8 | 44.5 | |
| SOUTH COAST | 18.6 | 16.8 | 17.0 | 9.2 | 7.5 | 7.1 | 79.0 | 73.7 | 50.3 | |
| EAST COAST** | 18.0 | 16.3 | 16.5 | 8.3 | 6.6 | 6.3 | 76.8 | 67.3 | 50.7 | |

^{*} West Coast and Inland Values cover the period 1983-2010

^{**} East Coast Temperature Values cover the period 1981-2007

Evaluation of November's seasonal forecast for the area of Cyprus

The seasonal forecast for **November** suggested that temperature of the island would be generally normal to above normal, over the eastern part, such as the mean monthly temperature of the surrounding area. The amount of the accumulated precipitation was expected to range generally from 50% to 75% of normal while over south-eastern area was expected to be below 50% of normal. Parts of the coastal zone of the surrounding area would receive again high precipitation accumulation.



Divergence of the mean monthly temperature (°C) from normal during November

Percentage of the mean monthly precipitation (%) compared with the normal of November

From the provisional data recorded by the Department of Meteorology, for the selected portion of the meteorological stations, which are presented in the table below,

| St. No. | Station Name | Mean Daily Maximum Temperature (°C) | Normal Value (1981- 2010) | Difference from Normal Value | Highest Daily Maximum Temperature (°C) | Lowest Daily Maximum Temperature (°C) | Mean Daily Minimum Temperature (°C) | Normal Value (1981- 2010) | Difference from Normal Value | Lowest Daily Minimum Temperature (°C) | Highest Daily Minimum Temperature (°C) | Monthly Total Precipitation (mm) | Normal Value (1981- 2010) | Difference from Normal Value |
|---------|------------------------|---|------------------------------------|---------------------------------------|---|--|--|------------------------------------|---------------------------------------|--|---|--|------------------------------------|---------------------------------------|
| 41 | POLIS CHRYSOCHOUS | 23.3 | 21.7 | 1.6 | 27.7 | 16.6 | 13.3 | 12.2 | 1.1 | 9.7 | 16.5 | 74.4 | 58.6 | 15.8 |
| 82* | PAFOS (AIRPORT) | 24.2 | 22.4 | 1.8 | 28.1 | 16.1 | 14.1 | 12.9 | 1.2 | 11.1 | 17.0 | 85.1 | 52.6 | 32.5 |
| 225 | PRODROMOS (C.F.C.) | 13.8 | 12.8 | 1.0 | 21.8 | 7.0 | 5.8 | 6.1 | -0.3 | 3.0 | 8.5 | 58.0 | 93.8 | -35.8 |
| 666* | ATHALASSA (RADIOSONDE) | 24.0 | 22.1 | 1.9 | 29.4 | 18.0 | 11.6 | 10.4 | 1.2 | 6.8 | 15.1 | 38.3 | 43.2 | -4.9 |
| 731 | LARNAKA (AIRPORT) | 24.2 | 22.6 | 1.6 | 27.3 | 19.1 | 15.3 | 12.4 | 2.9 | 12.0 | 19.3 | 27.0 | 46.8 | -19.8 |
| 800** | ACHNA (DASAKI) | 23.4 | 22.4 | 1.0 | 27.3 | 17.5 | 13.8 | 11.8 | 2.0 | 11.4 | 17.6 | 53.0 | 44.0 | 9.0 |
| | | * | | | | | Values cover es cover the | | | | | | | |

is evident that the model did not perform well concerning the temperature anomalies, since the most of both mean daily maximum temperatures and mean daily minimum temperatures were recorded above normal.

Extremes were recorded with positive departures greater than 4°C, like Prodromos station where the highest daily maximum temperature (21.8°C) was 9°C greater than normal (12.8°C) and like Athalassa station, where the highest daily maximum temperature (29.4°C) was 7.3°C greater than normal (22.1°C).


Extremes were recorded with positive departures greater than 4°C, like Larnaka station where the highest daily minimum temperature (19.3°C) was 6.9°C greater than normal (12.4°C) and like Achna station, where the highest daily minimum temperature (17.6°C) was 5.8°C greater than normal (11.8°C).

As regarding the accumulated precipitation **November**, the accumulated precipitation was above normal, while again it wasn't well forecasted by the model. During the periods 3-21 and 25-30 of **November** local showers and isolated thunderstorms resulted in accumulated precipitation of 113% of normal.


It is worth mentioning that on the 9th, 11th and 15th of **November** hail was reported.

Also, for the 9th, 11^{th} , 12^{th} , 15^{th} , 19^{th} , 20^{th} , 21^{st} , 28^{th} $\kappa\alpha\iota$ 30^{th} of **November** EMMA yellow warning for showers and thunderstorms were issued.

For the purpose of better visualization of **November** accumulated precipitation, a chart of Cyprus with the total preliminary accumulated precipitation is presented.



A Cyprus chart with the normal (period 1981 to 2010) accumulated precipitation for the month of **November** is also presented.



(*) It is stated that due to the failure of the seasonal model to correctly forecast the expected precipitation (sometimes) the seasonal forecast for precipitation is given with a reserve.