Seasonal weather forecast for the months of June, July and August 2020


General over view of the weather expected to prevail for June, July and August.

The weather for the following three months, namely **June**, **July** and **August 2020**, will be presented, with respect to the anomalies of temperature, and accumulated precipitation.

Normal temperatures (both maximum and minimum), are on the rise suggesting the second half of the period as the hottest period of the year. The accumulated precipitation is at its lowest values. The period of the following three months is likely to be characterized by outbreaks of thunderstorms, initiated from thermal causes over the main plain area or over the mountainous range from orographic uplift.

Specifically, regarding the seasonal forecast for the period of **June**, **July** and **August 2020**, the temperature will be generally normal. That means that the mean temperature will be around normal and does not exclude days with temperatures well above normal. Normal temperature also characterizes the greater part of Asia Minor, while the Balkans will be warmer than normal and near East will be lower than normal.

Regarding the seasonal forecast, based on the climatology of the island, the accumulated precipitation is expected to be below normal (as shown in the tables below). Any precipitation in Cyprus will be a result of thermal instability, in the form of a thunderstorm, with very local and temporal characteristics. For the above reason, but also due to the inability of the seasonal model to estimate precipitation for the region of Cyprus due to thermal instability, no extended comment will be made.




Divergence of temperature from normal from the mean seasonal temperature (°C) for June, July and August

Percentage (%) of the mean seasonal accumulated precipitation compared to the mean normal for June, July and August

The seasonal monthly forecast in detail

June

The seasonal forecast suggests that **June's** temperature will be generally normal while the Balkans and Middle East will be warmer than normal. Regarding the accumulated precipitation Cyprus is expected to be dry without the occurrence of dynamic barometric weather systems.




Divergence of the mean monthly temperature ($^{\circ}$ C) from normal during June

Percentage of the mean monthly precipitation (%) compared with the normal of June

July

The seasonal forecast suggests that **July's** temperature will be again normal. The accumulated precipitation at west part of Cyprus is expected to be low but over the east part and inland the accumulated precipitation is expected to be normal and higher than normal, respectively. Bearing always in mind that the normal accumulation for July is very low especially for the main plain and coastal areas




Divergence of the mean monthly temperature (°C) from normal during July

Percentage of the mean monthly precipitation (%) compared with the normal of July

August

The seasonal forecast suggests that **August's** temperature will be generally normal. The accumulated precipitation is expected to be low, bearing again in mind that the normal accumulation for August is very low especially for the main plain areas.



Divergence of the mean monthly temperature (°C) from normal during August

Percentage of the mean monthly precipitation (%) compared with the normal of August

Normal values of temperature (mean maximum and mean minimum) and accumulated precipitation for June, July and August

The normal values of mean maximum mean minimum temperature and accumulated precipitation are presented below concerning the three months the period of forecast is covering, in order to gain a better view of the normal seasonal climate. The period of forecast is the transitional summer. The temperature, both maximum and minimum, is on the rise with July's temperature significantly higher. Climatologically extreme high temperatures are recorded during July and over some areas during August. On the other hand, the accumulated precipitation is declining, as presented by the climatological data shown on the table below. Thundery activity is likely during afternoon hours mainly inland and over the mountains, which is noted also from the table below. Note the cause of the mountainous area of Troodos range over the accumulated precipitation, mainly because of thundery activity, which is affecting primarily the mountainous ranges of the island. The accumulated precipitation over Prodromos station although declining towards August is always significantly higher, if compared to the accumulated precipitation of the other selected stations.

TEMPERATURE AND PRECIPITATION NORMAL VALUES FOR THE PERIOD 1981-2010									
	MEAN DAILY MAXIMUM TEMPERATURE (°C)			MEAN DAILY MINIMUM TEMPERATURE (°C)			MEAN MONTHLY TOTAL PRECIPITATION (mm)		
Area Name	June	July	August	June	July	August	June	July	Auguist
NORTH COAST	30.3	33.4	33.3	18.5	21.1	21.5	1.8	0.1	0.0
WEST COAST-	27.6	29.9	30.5	18.0	20.6	21.1	1.3	0.2	0.0
MOUNTAINOUS AREAS	25.0	27.9	28.0	15.0	18.2	18.1	27.2	16.4	12.0
INLAND:	34.0	37.1	36.9	19.1	22.1	21.9	11.6	4.2	1.8
SOUTH COAST	30.2	32.5	32.9	19.4	22.0	22.3	2.0	0.5	0.3
EAST COAST**	31.0	33.2	33.5	18.9	21.8	22.0	1.8	0.1	0.1


^{*} West Coast and Inland Values cover the period 1983-2010

Evaluation of Mays seasonal forecast for the area of Cyprus

May

The seasonal forecast suggested that **May's** temperature would be generally normal with the western part being possibly warmer by 1° to 2°C than normal, while normal to above normal was suggested to be over most parts of the surrounding area. Regarding the accumulated precipitation was expected to range below 50% of normal over the island.

^{**} East Coast Temperature Values cover the period 1981-2007



Divergence of the mean monthly temperature (°C) from normal during May

Percentage of the mean monthly precipitation (%) compared with the normal of May

Concerning the models' evaluation for **May**, a general judgment is that it did not perform well neither regarding the temperature, either regarding the accumulated precipitation.

In a more detailed evaluation of the recorded temperatures both mean daily maximum and minimum (table below) were above normal. As an example, is Pafos airport with a month's absolute maximum of 42.5°C and Polis Chrysochous station with 41.7°C, with the normal being respectively 24.5°C and 25.8°C. Extreme minimums with positive departure from normal were also recorded like at Larnaka airport highest minimum of 29.5°C and Athalassa 27.5°C with the normal being respectively 15.7°C and 14.8°C.

From the distribution (provisional accumulated precipitation chart) of the accumulated precipitation of the month the mean surface distribution was near normal, reaching 14.3 mm or 89% of normal.

The days with precipitation were the 1st, 7th, 9th and 12th of **May**.

For the days 1, 2, 4, 5, 8, 9, 20, 22, 23 and 28 of **May** local showers, accompanied with thunderstorms sometimes, were recorded. It is worth mentioning that on the 5th and 23rd of **May** hail was reported.

On the 5th of **May** EMMA yellow level warning was issued concerning thunderstorms and on the 16th, 17th, 18th, 19th, 20th and 21st of **May** EMMA yellow and orange level warnings were issued concerning high temperatures.

TEMPERATURE AND PRECIPITATION PROVISIONAL DATA FOR MAY 2020

St. No.	Station Name	Mean Daily Maximum Temperature (°C)	Normal Value (1981- 2010)	Difference from Normal Value	Highest Daily Maximum Temperature (°C)	Lowest Daily Maximum Temperature (°C)	
41	POLIS CHRYSOCHOUS	27,4	25,8	1,6	41,7	20,8	
82*	PAFOS (AIRPORT)	26,8	24,5	2,3	42,5	22,1	
225	PRODROMOS (C.F.C.)	22,8	20,4	2,4	33,2	12,0	
666*	ATHALASSA (RADIOSONDE)	31,7	29,6	2,1	43,9	23,1	
731	LARNAKA (AIRPORT)	29,1	26,4	2,7	40,6	23,4	
800**	ACHNA (DASAKI)	29,1	27,4	1,7	42,8	21,3	

Mean Daily Minimum Temperature (°C)	Normal Value (1981- 2010)	Difference from Normal Value	Lowest Daily Minimum Temperature (°C)	Highest Daily Minimum Temperature (°C)
15,7	14,3	1,4	11,9	24,1
16,0	14,7	1,3	11,5	24,6
11,3	10,8	0,5	3,8	21,2
16,1	14,8	1,3	10,6	27,5
18,3	15,7	2,6	12,9	29,5
16,5	14,9	1,6	11,3	27,0


Monthly Total Precipitation (mm)	Normal Value (1981- 2010)	Difference from Normal Value	
2,5	7,6	-5,1	
2,7	6,1	-3,4	
12,9	32,0	-19,1	
traces	24,6	-24,6	
6,8	9,8	-3,0	
11,2	10,7	0,5	

= VALUES FROM AUTOMATIC WEATHER STATION

 $[\]ensuremath{^{*}}$ Pafos' and Athalassa's Station Normal Values cover the period 1983-2010

^{**} Achna's Temperature Normal Values cover the period 1981-2007

For the purpose of better visualization of **May** accumulated precipitation, a chart of Cyprus with the total preliminary accumulated precipitation is presented.



A Cyprus chart with the normal (period 1981 to 2010) accumulated precipitation for the month of **May** is also presented.

